

PERMANENT COSMETICS AND MICROBLADING: Possible Risks, Hazards, or Complications

Pain: There can be pain even after the topical anesthetic has been used. We do everything in our power to make the client as comfortable as possible, anesthetics work better on some people than on others.

Infection: Infection is very unusual and extremely rare. The areas treated must be kept clean, and only freshly cleaned hands should touch the treated areas. See "After Care" sheet for instructions on how to take care of treated area.

Uneven Pigmentation: This can result from poor healing, infection, bleeding or lifestyle after treatment. Your follow-up appointment will likely correct any uneven appearance.

Asymmetry: Every effort will be made to avoid asymmetry, but our faces are not symmetrical, so adjustments may be needed during the follow-up session to correct any unevenness.

Excessive Swelling or Bruising: Some people bruise or swell more than others. Ice packs ay help reduce the swelling. The swelling or bruising typically disappears between 1-2 days. Most people do not bruise at all and very few have swelling past a few hours post treatment.

Anesthetics: Topical anesthetics are used to numb the area to be treated. Lidocaine, Prilocaine, Benzocaine, Tetracaine, and/or Epinephrine cream and/or Gel are used. If you are allergic to any of these, please inform us.

MRI: Because pigments used in Permanent Cosmetic procedures contain inert oxides, a low-level magnet may be required if you need to be scanned by an MRI machine. You must inform your MRI Technician of any tattoos or permanent cosmetics that you have had.

If you have any other questions beyond your consultation about possible hazards associated with your procedure please let us know.

By signing below, you are agreeing that you have read and understand the possible risks and hazards associated with your procedure and have had all questions answered in a satisfactory manner.

Consent and Release for procedures performed:

Signed: _____ Date: _____